

# Smoking estimates from around the world: data from the first 17 participating countries in the World Mental Health Survey Consortium

Carla L Storr, Hui Cheng, Jordi Alonso, et al.

*Tob Control* 2010 19: 65-74 originally published online December 4, 2009 doi: 10.1136/tc.2009.032474

Updated information and services can be found at: http://tobaccocontrol.bmj.com/content/19/1/65.full.html

These include:

References

This article cites 34 articles, 19 of which can be accessed free at: http://tobaccocontrol.bmj.com/content/19/1/65.full.html#ref-list-1

n Red

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

**Notes** 

To order reprints of this article go to: http://tobaccocontrol.bmj.com/cgi/reprintform

# Smoking estimates from around the world: data from the first 17 participating countries in the World Mental Health Survey Consortium

Carla L Storr, <sup>1,2</sup> Hui Cheng, <sup>3</sup> Jordi Alonso, <sup>4</sup> Matthias Angermeyer, <sup>5</sup> Ronny Bruffaerts, <sup>6</sup> Giovanni de Girolamo, <sup>7</sup> Ron de Graaf, <sup>8</sup> Oye Gureje, <sup>9</sup> Elie G Karam, <sup>10,11</sup> Stanislav Kostyuchenko, <sup>12</sup> Sing Lee, <sup>13</sup> Jean-Pierre Lepine, <sup>14</sup> Maria Elena Medina Mora, <sup>15</sup> Landon Myer, <sup>16</sup> Yehuda Neumark, <sup>17</sup> Jose Posada-Villa, <sup>18</sup> Makoto Watanabe, <sup>19</sup> J Elisabeth Wells, <sup>20</sup> Ronald C Kessler, <sup>21</sup> James C Anthony<sup>3</sup>

For numbered affiliations see end of article.

#### Correspondence to

Dr Carla Storr, 655 W Lombard Street, Suite 655A, Baltimore, MD 21201, USA; cstor002@ son.umaryland.edu

Received 30 June 2009 Accepted 29 October 2009 Published Online First 4 December 2009

#### **ABSTRACT**

**Objective** To contribute new multinational findings on basic descriptive features of smoking and cessation, based upon standardised community surveys of adults residing in seven low-income and middle-income countries and 10 higher-income countries from all regions of the world.

**Methods** Data were collected using standardised interviews and community probability sample survey methods conducted as part of the WHO World Mental Health Surveys Initiative. Demographic and socioeconomic correlates of smoking are studied using cross-tabulation and logistic regression approaches. Within-country sample weights were applied with variance estimation appropriate for complex sample survey designs.

**Results** Estimated prevalence of smoking experience (history of ever smoking) and current smoking varied across the countries under study. In all but four countries, one out of every four adults currently smoked. In higher-income countries, estimated proportions of former smokers (those who had quit) were roughly double the corresponding estimates for most low-income and middle-income countries. Characteristics of smokers varied within individual countries, and in relation to the World Bank's low-medium-high gradient of economic development. In stark contrast to a sturdy male-female difference in the uptake of smoking seen in each country, there is no consistent sex-associated pattern in the odds of remaining a smoker (versus quitting).

**Conclusion** The World Mental Health Surveys estimates complement existing global tobacco monitoring efforts. The observed global diversity of associations with smoking and smoking cessation underscore reasons for implementation of the Framework Convention on Tobacco Control provisions and prompt local adaptation of prevention and control interventions.

#### INTRODUCTION

In recent years, there have been major advances in our understanding of the descriptive epidemiology of tobacco smoking, other forms of tobacco consumption and tobacco attributed mortality and morbidity. <sup>1–7</sup> An estimated quarter to a third of the world's population aged 15 years and older consumes tobacco on a daily basis. <sup>8</sup> Surveillance and monitoring of smoking activity on a global level are

important public health tasks in the context of effective tobacco control efforts.

Surveillance and research accomplishments during the 1990s provided a glimpse of national differences of tobacco involvement. The WHO assembled information from surveys and administrative records so to permit publication of the 2003 edition of the WHO Tobacco Control Country Profiles. These profiles characterise the tobacco situation in 196 countries and territories around the world.<sup>3</sup> Nevertheless, careful study of betweencountry variations in the tobacco smoking survey methods prompt cautious interpretation; many samples were non-representative; study designs and smoking assessments varied substantially. In another important cross-national study on the topic of nicotine prevalence and dependence, Fagerstrom and colleagues constrained some of the sources of variation and were able to present estimates based solely upon standardised nicotine dependence assessments. Nonetheless, even here there was some unevenness in methods of sampling and contexts for assessment (eg, in the clinical samples). 10 11

Problems of this type can be overcome when research teams from different countries agree to follow a common protocol, as has been done in school surveys of tobacco and other drug involvement of young people in different regions of the world. <sup>2 4 12 13</sup> Standardised surveys from probability samples of adults from numerous countries are also beginning to guide global tobacco control efforts.9 14-16 In addition, the first ever international cohort study of tobacco use, the International Tobacco Control Policy Evaluation Project, is now conducting parallel prospective surveys of representative samples of adult smokers in at least 20 countries (inhabited by over 50% of the world's population, 60% of the world's smokers), in an effort to evaluate the psychosocial and behavioural impact of the Framework Convention on Tobacco Control (FCTC). 17 18

In the background of these investigations, there is a tobacco epidemic model synthesised by Lopez and colleagues, on the basis of previous surveillance data and several indicators of disease burden. <sup>19</sup> The various epidemic stages described in this model highlight overall differences in the prevalence of smoking and attributable disease and deaths, as well as characteristic differences among users, such as

ratios of male to female smokers. In many low-income and middle-income countries (generally characterised as Stage 1 and Stage 2 in the Lopez model), the estimated prevalence and consumption of tobacco cigarettes have been rising steadily. 9 20 There is hope that vigorous prevention and early intervention efforts may help some of the low-income and middle-income countries avoid transitions to more advanced stages of increased consumption and disease burden that characterise later model stages. For example, many countries now are engaged in preventive actions to dissuade youths from smoking uptake. Nonetheless, each country will also need brief interventions and other aids to encourage quitting among smokers. Smokers who quit thereby limit the number of years of smoking and gain substantial benefits to health, wellbeing, and increased survivorship. <sup>21–23</sup> In some high-income countries there have been recent increases in the percentage of former smokers, but generally little change in the percentage of former smokers has been found in low-income and middle-income countries where population-level monitoring has occurred.8

In consequence, along with surveillance of smoking uptake and associated health indicators, a population survey focus on former smokers, smoking cessation and characteristics associated with smoking cessation should be fruitful, providing additional clues about the course of the global tobacco epidemic. For this reason, our research group was granted permission to produce estimates of smoking prevalence, as well as to study former smokers, drawing upon some recently gathered data about mental health characteristics in 17 different countries from all regions of the world, representing a range of the higherincome, middle-income and low-income jurisdictions, during the first wave of the WHO World Mental Health Surveys Initiative (WMHS, http://www.hcp.med.harvard.edu/wmh). In this paper, we are able to characterise more than the 'depth' of the smoking epidemic in each country (as manifest in the countryspecific prevalence estimates); we also gain a perspective on each country's progress in the promotion of smoking cessation in various population subgroups within each country—for example, as manifest in male-female differences in who has quit versus who has continued to smoked. For each regional site or country as a whole, and for males and females considered separately, the report provides estimates of smoking, based on cross-sectional survey data, with due attention to a two-group classification of smokers—that is, current smokers versus former or ex-smokers. Prevalence estimates are given for selected correlates of current smoking status (eg, age group) so as to help build up a more solid foundation of empirical survey findings, upon which tobacco researchers might construct more detailed and probing studies of the processes and conditions that foster smoking uptake and cessation in each place.

#### **METHODS**

# Populations and sampling approach

The WMHS Initiative continues to unfold, with new countries participating in successive waves, as resources become available and survey research teams learn and implement the standardised field survey protocol. This report is based upon WMHS data from the first 17 participating countries, with sites spread throughout all the major regions of the world. The consortium has a mental health focus, and its main goal is to estimate prevalence and impact of selected psychiatric disorders such as major depression and the anxiety disorders, worldwide, via standardised interviews and a common survey protocol. <sup>24–26</sup> The countries, grouped by major region, encompass a range of

economies as classified by the World Bank<sup>27</sup>:the Americas (Colombia, Mexico and the USA), Asia/Pacific (People's Republic of China (PRC), Japan and New Zealand), Europe (Belgium, France, Germany, Italy, The Netherlands, Spain and the Ukraine), and the Middle East and Africa (Israel, Lebanon, Nigeria, South Africa). While in many of the countries it was feasible to assess a nationally representative sample, estimates for several countries are based on area samples (table 1). The sample for Nigeria approximated 57% of the population, samples from Colombia and Mexico approximated 73% of their populations and samples for Japan and the PRC were selected from large metropolitan areas.

In each jurisdiction, nationally (or regional) representative household samples of adult residents were obtained via a multistage probability sample survey approach. To date, a total of over 80 000 adults have completed WMHS assessments, with participation levels as shown in table 1. Estimates reported in this paper are for adults age 18 years and older at the time of assessment; younger teenagers were surveyed in New Zealand and South Africa, where the research teams have prepared separate reports on smoking and other health characteristics of 15–17 year olds as part of internal country reports on their site-specific projects. In most countries, persons not speaking the country's primary languages were not sampled.

#### Field procedures

Training and field procedures for sampling, recruitment and assessment were standardised across countries. 24 26 In almost all countries, trained lay interviewers carried out face-to-face recruitment and consent procedures, followed by either paper and pencil or computerised assessments. Informed consent was obtained after each individual had received a description of the study goals and procedures, data uses and protections, and participant rights. The institutional review board of the organisation coordinating the survey in each country approved and monitored compliance in human subject protection and informed consent processes.

#### Measures

The interview was divided into two parts to reduce respondent burden. Data for this paper are from the smoking screening questions assessed in the first part, completed by all participants at every site. This first part included standardised items on personal characteristics such as age, sex, marital status, a set of health behaviour screening items and a core diagnostic assessment for selected psychiatric disturbances. Items on employment and education were optional elements of this 'Part 1' assessment, and were not assessed in every country.

The key response variable in this report, for all countries except Israel, is based upon a self-classification of respondents who answered this question "Are you a current smoker [active], ex-smoker [former], or have you never smoked?" The Israel survey did not have the same question. Instead smoking status was identified by a set of responses to these two questions: "Are you currently smoking at least once a day [active]?" and "Have you ever smoked in the past, at least once a day [former]?". Some sites also added a more comprehensive tobacco module in the second part of the interview with specifications for how much, how often, age of onset, etc (eg, cigarettes, cigars, pipes and other types of tobacco products), but the uneven availability of these data constrains multi-country analyses of the type reported here; these details are being reported by the individual sites in their own papers. <sup>28</sup>

Table 1 Characteristics of the first 17 population surveys for the World Mental Health (WMH) Survey Consortium

Site	No	Age range	Year fielded	Level of study participation (%)	Sample composition
Americas					
Colombia	4426	18-65	2003	88	Household residents in all urban areas of the country (approx 73% of total national population)
Mexico	5782	18-65	2001—2	77	Household residents in all urban areas of the country (approx 73% of total national population)
USA	9282	≥18	2002—3	71	Household residents, nationally representative
Asia and Pacific	Region				
Japan	1663	≥20	2002—3	56	Household residents in four metropolitan areas: Fukiage, Kushikino, Nagasaki, Oyayama
People's Republic of China (PRC)	5201	≥18	2002—3	75	Household residents in Beijing (2633) and Shanghai (2568) metropolitan areas
New Zealand	12790	≥18*	2003-4	73	Household residents, nationally representative
Europe					
Belgium	2419	≥18	2001—2	51	Households from national resident register, nationally representative
France	2894	≥18	2001—2	46	Households with working and listed telephone numbers, nationally representative
Germany	3555	≥18	2002—3	58	Individuals from community resident registries, nationally representative
Italy	4712	≥18	2001—2	71	Individuals from municipality resident registries, nationally representative
Netherlands	2372	≥18	2001—2	56	Household residents listed in municipal postal registries, nationally representative
Spain			2001—2		Household residents, nationally representative
Ukraine	4725	≥18	2002	78	Household residents speaking either Russian and Ukrainian, nationally representative
Middle East and	Africa				
Israel	4859	≥21	2002—4	73	Individuals from national resident register, nationally representative
Lebanon	2856	≥18	2002—3	70	Household residents, nationally representative
Nigeria	6752	≥18	2002—4	80	Household residents in 21 of the country's 36 states (approx 57% of total national population)
South Africa	4315	≥18*	2002—3	87	Household residents, nationally representative

<sup>\*</sup>New Zealand and South Africa samples restricted to those ≥18 years for this report.

Covariates of interest, also measured during interviews, included marital status categories, specified as currently married or living together, never married, or no longer married (widowed, separated or divorced) and five categories of education.

Employment status categories reflect each respondent's main activity at the time of the interview. When respondents were not working or were students, homemakers or retired, they were classified as 'other' (mainly unemployed people).

## Data analysis

Site-specific estimates are presented for proportions of smokers in relation to the characteristics just described. Estimation procedures for each country took into account differential probabilities of selection within households and communities, as well as post-stratification balance to known population census distributions (eg, age, sex). All estimates shown are based upon within-country weights and post-stratification adjustment, without a shift to world standard population distributions. In subsequent steps, we used simple cross-tabulations to study variations in smoking status (never vs ever and former versus current). Covariate-adjusted OR estimates were estimated via the generalised linear model with a logistic link to convey the strength of associations that link being male or female with being an ever-smoker and with respect to the active/former smoking contrast. Appropriate for complex sample survey designs, the survey commands and the Taylor series linearisation approach in Stata versions 9-10 were used for variance estimation and to estimate p values.<sup>29</sup>

#### **RESULTS**


#### **Country-specific smoking estimates**

Estimated prevalence of smoking experience varied across the countries under study (figure 1). Typically, the estimated prevalence of ever-smoking (ie, active and former smokers combined) was lower in low-income and middle-income countries (14%—47%) compared to estimates for high-income countries (range 47%—60%). Throughout most of Europe, the USA, Israel and Japan, almost one-half of adults in the sampled community population initiated smoking; either they continue to smoke currently or they had smoked at sometime in the past. This estimated cumulative incidence proportion was largest in The Netherlands, where nearly three out of five adults had started smoking at some point during their lifetime. At the other extreme is Nigeria, with an estimate under 20%.

We also see cross-national variation in the estimates derived from the WMHS cross-sectional survey data on current tobacco smoking (table 2). For example, Nigeria again provides an extreme value. In that country, the estimated fraction of current smokers among all adults is less than 1 in 20 (point prevalence=3.9). Columbia also shows a relatively low estimate at 14.6%. The estimated prevalence of current smoking is between 20% and 30% in nine of the countries studied, and in six countries the current smoking point prevalence estimates exceed 30%. Countries from each of the three income tiers are represented in the high prevalence stratum (Lebanon, Spain, Ukraine, PRC (Beijing and Shanghai sites), The Netherlands and Germany). As for being in the lowest rank with respect to estimated prevalence of current smoking among high-income countries, New Zealand, the USA and Belgium share this distinction.

Viewed cross-nationally, there is also considerable variation in estimates for former or ex-smoking (table 2). For example, in Nigeria, about three in every four smokers has quit; at the other extreme in the low-income and middle-income countries, for Lebanon, South Africa and China, the great majority of eversmokers continue to smoke and did not qualify as former smokers. In European countries, roughly one-third to one-half of the smokers had quit. At the other extreme, in the Americas

Figure 1 Estimated overall prevalence of being an active or former smoker, by country, as well as the prevalence for never smoking. World Mental Health Survey Consortium Surveys.



# \* World Bank denoted low or middle income country

(Colombia, Mexico, USA), and in New Zealand, more than one-half had quit, describing themselves as former smokers.

# Estimated proportion of smokers by selected background characteristics

Across countries, there is some substantial population subgroup variation in the estimated prevalence of current smoking, as can be seen in table 3 (low-income and middle-income countries) and table 4 (higher-income countries). Against a background of a generally sturdy male excess prevalence of smoking, relative to estimates for females, the male excess was rather modest or non-existent in New Zealand, Germany, The Netherlands and the USA, where roughly one-fifth to one-third of men and women qualified as current smokers. In contrast, nearly all adult current

smokers in China and Nigeria were men; in South Africa, Ukraine and Japan, current smoking also was substantially more common among men.

With respect to youngest adult age group (18–24 year olds), Germany and Ukraine had current smoking prevalence estimates above 45% (almost 1:1 odds of smoking), compared to other countries with corresponding values as small as one current smoker per 100 18–24 year olds in Nigeria (1.2%) and one per eight in Colombia (11.9%), and as large as one in 2.5 in Spain (42.1%). Current smoking prevalence among 18–24 year olds living in most of the high-income countries is often as high or higher than that for 25–44 year olds—something not seen among the low/middle-income countries. Current smoking prevalence estimates above 45% were observed only for

**Table 2** Estimated overall prevalence (95% CI) of smoking status by country:WMH survey consortium, 2001—2005

	Current/active	Former/quit	Never smoked
Lebanon*	36.0 (33.2 to 12.3)	10.5 (8.7 to 12.3)	53.4 (50.7 to 56.1)
Spain	34.2 (32.7 to 35.7)	18.4 (16.8 to 19.9)	47.4 (45.6 to 49.1)
Ukraine*	32.4 (30.8 to 34.0)	13.3 (12.1 to 14.5)	54.2 (52.6 to 55.9)
China*	31.8 (29.0 to 34.8)	6.9 (5.6 to 8.5)	61.3 (57.6 to 64.4)
Netherlands	31.8 (29.7 to 33.7)	27.6 (25.2 to 30.0)	40.6 (38.1 to 43.1)
Germany	31.2 (29.4 to 33.0)	20.0 (17.7 to 22.2)	48.8 (46.6 to 51.0)
France	27.6 (25.6 to 29.7)	24.1 (22.1 to 26.1)	48.2 (45.2 to 51.3)
Israel†	27.5 (26.3 to 28.8)	20.1 (18.9 to 21.3)	52.4 (51.0 to 53.8)
Italy	27.4 (25.8 to 29.1)	20.0 (18.7 to 21.4)	52.5 (50.4 to 54.6)
Japan	27.0 (25.1 to 28.9)	22.7 (20.9 to 24.4)	50.3 (48.1 to 52.3)
Belgium	25.7 (23.5 to 27.8)	23.2 (20.7 to 25.7)	51.2 (48.6 to 53.7)
South Africa*	25.6 (23.9 to 27.2)	9.6 (8.3 to 10.9)	64.8 (62.8 to 66.6)
USA	25.0 (23.8 to 26.1)	27.6 (26.4 to 28.8)	47.3 (45.8 to 48.9)
New Zealand	23.4 (22.4 to 24.4)	27.8 (26.8 to 28.8)	48.7 (47.4 to 49.9)
Mexico*	20.6 (18.8 to 22.4)	27.1 (25.3 to 28.9)	52.2 (50.0 to 54.4)
Colombia*	14.6 (13.2 to 15.9)	23.9 (22.2 to 25.5)	61.5 (59.7 to 63.3)
Nigeria*	3.9 (3.5 to 4.3)	10.4 (9.4 to 11.4)	85.7 (84.6 to 86.8)

<sup>\*</sup>Denotes low/middle-income countries as classified by the World Bank. †Israel estimates are for those who smoked at least once a day.

25–44 year olds in Spain and 35–54 year olds in Lebanon. Uniformly, current smoking prevalence estimates were lower for adults age 55 years and older, compared to their younger adult counterparts, most likely due in part to smoking-attributable excess mortality for smokers, although the contrast between 45–54 year olds and those 55 years and older was not especially marked in Mexico, Italy, Israel, The Netherlands and the USA.

The education gradients vary substantially across countries—in many countries, including the USA, prevalence is lower

among those with no or some primary education than it is for those who finished primary schooling. In general, the highest current smoking prevalence estimates are observed for adults whose highest educational attainment was at the primary or secondary education levels. In many but not all of the countries, the proportion of smokers with minimal education was quite similar to the proportion of smokers found among those with the most education (ie, college graduates). In some countries smoking appears to be less common among college graduates while in other countries college graduates are more likely to

Table 3 Estimated prevalence of current smoking by selected sociodemographic characteristics in each of the low-income to middle-income countries (%. SE): World Mental Health (WMH) Survey Consortium

	Nigeria (n = 282)	Peoples' Republic of China (n = 1584)	South Africa (n = 1010)	Colombia (n = 669)	Mexico (n = 1122)	Ukraine (n = 1285)	Lebanon (n = 1113)
Sex							
male	7.5 (0.5)	55.5 (1.3)	41.4 (1.3)	20.2 (1.3)	31.7 (1.4)	57.7 (1.3)	42.4 (1.8)
female	0.4 (0.2)	5.5 (0.6)	11.8 (0.9)	9.9 (0.7)	10.5 (0.8)	11.7 (0.6)	29.8 (1.8)
Age at interview (years)*							
18-24	1.2 (0.3)	16.9 (1.8)	21.4 (1.4)	11.9 (1.2)	19.4 (1.5)	45.6 (2.2)	24.5 (2.5)
25-34	4.6 (0.5)	37.9 (2.8)	28.6 (1.6)	11.6 (1.1)	20.2 (1.6)	47.9 (2.4)	32.8 (2.1)
35-44	5.3 (0.7)	40.4 (1.3)	30.2 (2.4)	17.4 (1.7)	24.2 (1.5)	42.2 (1.8)	49.2 (2.4)
45-54	6.1 (1.0)	36.1 (1.8)	25.0 (2.0)	19.5 (1.7)	21.8 (2.2)	32.3 (1.7)	48.5 (2.3)
55 and older	3.6 (0.7)	21.0 (1.3)	20.6 (1.9)	13.1 (1.7)	15.4 (1.8)	14.4 (0.8)	32.7 (3.0)
Education*							
None/some primary	2.6 (0.4)	Not assessed	25.8 (1.7)	21.0 (1.8)	18.8 (1.0)	9.5 (2.6)	37.5 (2.3)
Finished primary	5.3 (0.4)		26.2 (1.2)	14.7 (1.0)	22.2 (1.2)	25.3 (1.2)	45.4 (2.1)
Finished secondary	1.9 (0.5)		25.7 (2.1)	11.8 (1.3)	19.0 (1.8)	39.8 (1.2)	32.2 (2.9)
Some college	5.0 (0.8)		24.0 (2.3)	14.2 (1.7)	23.1 (2.6)	33.4 (1.7)	24.9 (3.1)
College graduate	2.2 (1.1)		20.9 (3.9)	9.6 (1.9)	16.6 (1.9)	28.0 (2.3)	24.7 (2.5)
Marital status							
Married/cohabiting	4.5 (0.4)	33.6 (1.0)	24.7 (1.2)	14.9 (1.0)	20.8 (0.9)	33.6 (0.9)	41.3 (1.8)
No longer married	3.1 (0.9)	33.6 (3.2)	34.5 (3.7)	16.0 (2.0)	18.9 (2.0)	20.3 (1.3)	38.2 (3.6)
Never married	2.9 (0.4)	26.8 (2.1)	25.2 (1.1)	13.4 (1.1)	20.5 (1.8)	45.6 (2.6)	27.6 (2.2)
Employment*							
Working	4.6 (0.3)	37.4 (1.3)	Not assessed	16.9 (1.0)	26.7 (1.1)	41.0 (1.3)	40.0 (1.9)
Student	1.2 (0.4)	5.0 (1.7)		9.8 (2.6)	16.2 (2.7)	35.9 (4.8)	14.7 (3.1)
Homemaker	0.3 (0.3)	7.3 (3.2)		8.3 (0.9)	9.4 (0.9)	21.4 (2.7)	34.3 (2.2)
Retired	6.4 (2.4)	17.2 (0.9)		10.0 (2.5)	14.3 (3.6)	13.8 (0.7)	42.5 (4.2)
Other	3.7 (1.0)	44.0 (2.6)		19.5 (2.4)	30.5 (4.8)	45.5 (2.9)	38.6 (3.9)

<sup>\*</sup>Reflect values at the time of interview

Table 4 Estimated prevalence of current smoking by selected sociodemographic characteristics in each of the high-income countries (%, SE); World Mental Health (WMH) Survey Consortium

					New		Nether-			
	Japan (n = 622)	Spain (n = 1727)	Italy (n = 1316)	Israel* (n = 1330)	Zealand (n = 3566)	France (n = 802)	lands (n = 771)	Germany (n = 1121)	Belgium (n = 645)	USA (n=2454)
Sex										
male	45.5 (1.5)	40.6 (1.4)	33.6 (1.0)	37.1 (1.0)	24.5 (0.8)	34.4 (1.6)	32.7 (1.5)	34.4 (1.4)	32.8 (1.5)	27.7 (0.6)
female	11.1 (0.7)	28.4 (1.0)	21.8 (1.2)	18.6 (0.8)	22.4 (0.7)	21.5 (1.2)	30.9 (1.4)	28.3 (1.2)	18.8 (1.1)	22.6 (0.7)
Age at interview (years)	†									
18-24	39.8 (5.1)	42.1 (2.0)	34.9 (2.8)	32.8 (2.1)	35.3 (1.6)	39.9 (4.0)	33.3 (4.1)	45.1 (3.6)	41.6 (3.7)	29.3 (1.6)
25-34	33.2 (2.9)	48.1 (1.9)	34.6 (1.6)	32.9 (1.4)	31.0 (1.1)	40.3 (2.9)	34.1 (2.8)	40.2 (2.5)	24.6 (2.7)	28.9 (1.3)
35-44	37.6 (2.8)	47.9 (1.4)	35.3 (1.5)	34.1 (1.6)	23.7 (1.0)	35.8 (2.2)	35.4 (2.2)	42.5 (2.0)	29.4 (1.7)	29.8 (1.1)
45-54	32.8 (2.4)	34.0 (2.2)	32.0 (1.5)	29.4 (1.5)	22.4 (1.1)	26.1 (1.8)	37.3 (2.0)	37.6 (2.9)	33.0 (3.0)	24.1 (1.4)
55 and older	16.9 (1.1)	15.5(1.0)	15.4 (1.0)	14.9 (1.0)	12.9 (0.7)	11.9 (1.3)	24.2 (1.4)	14.3 (1.2)	15.0 (1.4)	17.4 (0.8)
Education†										
None/some primary	21.1 (18.7)	21.3 (1.3)	18.3 (1.1)	15.8 (2.0)	25.9 (6.7)	Not assessed	26.0 (3.9)	20.4 (11.5)	24.9 (2.6)	24.5 (3.8)
Finished primary	20.8 (1.3)	39.0 (1.7)	32.7 (1.3)	34.3 (1.8)	30.3 (1.0)		34.5 (2.3)	13.8 (4.6)	24.5 (1.4)	37.9 (1.4)
Finished secondary	29.1 (1.6)	39.6 (3.9)	31.8 (1.6)	33.3 (1.2)	25.6 (0.9)		37.4 (3.4)	31.0 (1.0)	27.2 (2.7)	30.5 (1.1)
Some college	28.1 (2.4)	38.8 (2.3)	33.8 (2.1)	28.4 (1.6)	14.6 (1.0)		36.4 (2.2.)	38.8 (3.2)	27.6 (3.0)	23.2 (1.1)
College graduate	32.2 (2.8)	37.9 (2.0)	24.2 (2.2)	17.8 (1.1)	10.3 (0.8)		24.3 (1.5)	38.4 (6.5)	23.4 (2.5)	12.3 (0.8)
Marital status										
Married/cohabiting	27.5 (1.1)	32.2 (0.8)	25.8 (0.8)	24.9 (0.8)	19.8 (0.6)	26.2 (1.4)	30.9 (1.4)	28.4 (1.0)	23.4 (1.0)	21.6 (0.7)
No longer married	19.8 (1.9)	19.8 (2.1)	21.2 (3.1)	26.4 (1.7)	25.9 (1.2)	22.3 (3.0)	35.4 (3.8)	30.0 (1.8)	22.1 (2.1)	30.3 (1.3)
Never married	31.4 (3.3)	44.5 (2.0)	34.0 (1.8)	37.7 (1.7)	32.2 (1.1)	37.4 (2.9)	32.9 (3.3)	40.8 (1.7)	36.9 (3.3)	28.5 (1.1)
Employment†										
Working	34.6 (1.4)	45.5 (1.1)	33.4 (1.0)	32.0 (0.9)	24.1 (0.6)	33.9 (1.3)	35.0 (1.3)	37.9 (1.3)	30.2 (1.5)	25.8 (0.7)
Student	29.8 (9.6)	27.3 (3.6)	16.5 (3.6)	32.9 (4.4)	22.2 (1.9)	41.7 (8.4)	16.6 (4.4)	35.2 (5.3)	31.9 (6.7)	18.5 (2.9)
Homemaker	7.2 (0.9)	21.8 (1.7)	24.4 (2.1)	11.8 (2.1)	31.1 (1.5)	15.6 (3.3)	24.8 (2.5)	26.3 (3.1)	16.2 (2.8)	17.7 (1.5)
Retired	20.2 (2.5)	15.0 (1.1)	13.7 (1.0)	12.1 (1.2)	10.3 (0.8)	10.2 (1.3)	21.5 (2.1)	12.9 (1.5)	13.9 (1.6)	14.8 (1.0)
Other	28.7 (4.2)	40.1 (2.8)	31.1 (3.2)	30.4 (1.9)	43.6 (2.6)	37.5 (3.3)	46.0 (4.8)	47.3 (3.1)	34.8 (3.4)	43.1 (2.1)

<sup>\*</sup>Israel estimates are for those who smoked at least once a day.

smoke than those with no or some primary education only. The prevalence of current smoking for college graduates is comparable to that of adults with some college (but no college degree) in the Ukraine, Lebanon, Japan, Spain and Germany.

With respect to marital status, in many of the European countries, a larger proportion of never married adults were current smokers, compared to other categories. Nonetheless, estimated prevalence was not always the lowest among those married—for example, in Lebanon, a higher proportion of married than never married adults were current smokers.

In terms of employment, with two exceptions, Nigeria and Colombia, more than one-quarter of the working adults in other countries were current smokers. The prevalence range is wide, from 24% of workers in New Zealand to 45% of workers in Spain. Current smoking among adult students varied from as low as 1–5% in Nigeria and China to above 30% among students in Ukraine, France, Belgium and Germany, as well as in Israel (where the estimates pertain to current daily smoking as explained in the Methods section). For the most part, as expected, the proportion of current smokers among homemakers reflected that among females in general, and the proportion among those retired reflected the prevalence in the oldest age group. In all countries but one, the 'other' employment category (consisting largely of unemployed adults) was a subgroup with quite high prevalence of current smoking, the exception was Nigeria.

## Male-female differences

Table 5 has a focus on male-female variation in the occurrence of smoking, both past and current. In all but one country (Nigeria), more than 50% of the men had a history of having smoked (up to approximately 80% in Ukraine and Japan). Lower values were

observed for women such that there is a tangible association between being male and a history of smoking in all countries under study (including Nigeria; see first columns of table 5). A history of smoking tended to be more common among females in the high-income countries (in particular, nearly 50% in New Zealand, The Netherlands and the USA). In every country a male-female variation in the odds of ever smoking was noted (table 5). In Belgium, France, Italy, Spain, Mexico, Israel and Colombia, the odds of ever smoking among males were three times greater than the corresponding odds of smoking among females in these countries (adjusted OR ranged from 2.9–3.7). Especially large OR estimates were found in Japan, the Ukraine, South Africa, China and Nigeria, owing largely to generally smaller smoking proportions for the women in these countries.

In stark contrast to the sturdily replicated male-female variations in the uptake of smoking seen across these countries, there is no similar sturdy cross-national pattern in the prevalence or odds of remaining a smoker (versus quitting) for males versus females (table 5). In the higher-income countries, roughly 30%-50% of ever-smokers of both sexes have stopped smoking; Spain is the exception among these countries (table 5; final columns). Among lower-income and middle-income countries, a relatively smaller proportion of ever-smokers had quit in China and Lebanon (<25%), and a relatively larger proportion had quit in Nigeria and Colombia (>60%). Once they had started smoking, females were just as likely as males to continue smoking (ie, be a current smoker) in Nigeria, China, South Africa, Colombia, Lebanon, Italy, New Zealand and Germany. The only country showing a statistically robust female excess in the odds of continuing to smoke among adult smokers was Spain, where 75% of the adult female ever smokers continued to smoke, versus 59% of the adult male ever smokers (aOR=0.7; p<0.001);

<sup>†</sup>Reflect values at the time of interview.

Table 5 Estimated sex-specific distribution (%) of smoking status, with covariate-adjusted OR estimates for the association between being male and smoking status. Data from each country participating in the World Mental Health (WMH) Survey Consortium

	Estimated proportion of never and ever smokers			Estimated pro former and cu among ever-si	Adjusted + CD	
	Never	Ever	Adjusted† OR (95% CI)	Former	Current	Adjusted† OR (95% CI)
Nigeria			<u> </u>			
Female	98.3	1.7	1.0	75.8	24.2	1.0
Male	72.5	27.5	26.6 (18.5 to 38.5)**	72.7	27.3	1.1 (0.4 to 3.3)
People's Republic of		=1.10				(0 10 0.0)
Female	92.9	7.1	1.0	22.6	77.4	1.0
Male	32.9	67.1	35.2 (27.7 to 44.6)**	17.3	82.7	1.0 (0.6 to 1.6)
South Africa			, , , , ,			, , , , , , , , , , , , , , , , , , , ,
Female	83.0	17.0	1.0	30.1	69.9	1.0
Male	43.2	56.8	7.7 (6.2 to 9.5)**	26.5	73.5	1.1 (0.9 to 1.4)
Colombia			,			,
Female	72.5	27.5	1.0	64.1	35.9	1.0
Male	48.5	51.5	2.7 (2.2 to 3.2)**	60.8	39.2	1.0 (0.07 to 1.4)
Mexico	10.0	01.0	2.7 (2.2 to 0.2)	00.0	00.2	1.0 (0.07 to 1.1)
Female	69.5	30.5	1.0	65.4	34.6	1.0
Male	33.2	66.8	3.7 (3.0 to 4.4)**	52.5	47.5	1.6 (1.2 to 2.2)*
Ukraine	00.2	00.0	0.7 (0.0 to 4.4)	32.3	47.0	1.0 (1.2 to 2.2)
Female	81.5	18.5	1.0	36.5	63.5	1.0
Male	20.8	79.2	19.7 (16.0 to 24.3)**	27.1	72.9	2.5 (1.9 to 3.2)**
Lebanon	20.0	73.2	13.7 (10.0 to 24.3)	27.1	72.5	2.3 (1.3 to 3.2)
Female	61.8	38.2	1.0	22.1	77.9	1.0
Male	44.9	55.1	2.0 (1.5 to 2.8)**	23.1	76.9	1.0 (0.07 to 1.6)
	44.9	33.1	2.0 (1.5 to 2.6)	23.1	70.9	1.0 (0.07 to 1.0)
Japan	77.0	22.0	1.0	F1 0	40.0	1.0
Female	77.2	22.8	1.0	51.2	48.8	1.0
Male	19.0	81.0	15.2 (12.0 to 19.4)**	43.8	56.2	2.2 (1.6 to 3.0)**
Spain	00.0	07.7	4.0	04.0	75.0	1.0
Female	62.3	37.7	1.0	24.8	75.2	1.0
Male	31.1	68.9	3.2 (2.8 to 3.8)**	41.0	59.0	0.7 (0.5 to 0.9)*
Italy	00.0	20.0	4.0	05.4	04.0	1.0
Female	66.2	33.8	1.0	35.4	64.6	1.0
Male	37.6	62.4	3.2 (2.7 to 3.6)**	46.1	53.9	0.8 (0.7 to 1.1)
Israel‡						
Female	66.0	34.0	1.0	45.2	54.7	1.0
Male	37.7	62.3	3.1 (2.8 to 3.5)**	40.3	59.7	1.4 (1.1 to 1.7)*
New Zealand						
Female	51.8	48.2	1.0	53.4	46.5	1.0
Male	46.3	53.7	1.3 (1.2 to 1.4)**	54.4	45.6	1.1 (1.0 to 1.3)
France						
Female	60.9	39.1	1.0	45.0	55.0	1.0
Male	34.3	65.7	3.0 (2.5 to 3.6)**	47.6	52.4	1.2 (1.0 to 1.5)*
Netherlands						
Female	44.7	55.3	1.0	44.1	55.9	1.0
Male	36.3	63.7	1.4 (1.0 to 1.9)*	48.7	51.3	1.0 (0.7 to 1.2)
Germany						
Female	55.8	44.2	1.0	36.0	64.0	1.0
Male	41.3	58.7	1.7 (1.4 to 2.0)**	41.1	58.6	0.9 (0.7 to 1.3)
Belgium						
Female	63.5	36.5	1.0	48.4	51.6	1.0
Male	38.2	61.8	2.9 (2.5 to 3.4)**	46.8	53.2	1.4 (1.0 to 1.8)*
USA						
Female	52.4	47.6	1.0	52.6	47.4	1.0
Male	41.8	58.2	1.6 (1.5 to 1.8)**	52.4	47.6	1.1 (1.0 to 1.3)

a special opportunity for gender-targeted smoking outreach and intervention exists in that country.

In contrast with the pattern in Spain, a male excess in the odds of smoking persistence was found in six countries. In Ukraine and Japan, the evidence indicates that male smokers were more likely to continue smoking than their female smoking counterparts (aOR=2.5 and aOR=2.2, respectively, p<0.001). Adult male smokers in Israel, Mexico, France and Belgium are modestly more likely to remain active smokers, compared to female smokers (aOR=1.4, 1.6, 1.2 and 1.4 for Israel,

<sup>\*</sup>p<0.05, \*\*p<0.001.

<sup>†</sup>Adjusted holding constant age, marital status, education (except for China and France), and employment (except for South Africa).

<sup>‡</sup>Israel estimates are for those who smoked at least once a day.

Mexico, France and Belgium, respectively, p<0.05). It is noteworthy that in Israel, France and Belgium, somewhat more than one-half of ever-smokers were active smokers; this was not the case in Mexico where smokers were more likely to have quit than to have remained active smokers.

#### **DISCUSSION**

This study, with multiple sites around the world using common standardised research protocols, provides estimates to describe (1) adult smoking experiences and (2) patterns of association for continuing to smoke among smokers in relation to being a male or female. A few design and measurement issues merit attention before a detailed discussion of the findings. The sampling approach (eg, nationally representative vs specific areas within the country), eligibility requirements (eg, age range differences) and the items assessing smoking status were not entirely uniform across countries. Furthermore, as mentioned in the introduction, this research consortium was focused on mental disorders and the study did not focus exclusively on 'tobacco' smoking or the features of smoking as they vary from place to place (eg, use of water pipe in Lebanon versus commercial or hand-rolled cigarettes elsewhere). Reporting biases may occur as the willingness to identify oneself as a current or past smoker might vary across national boundaries or across the population subgroups under study.<sup>30</sup> In some countries, the mental health focus of the field research might have led some smokers to decide not to participate. Limitations such as these may affect these estimates to an unknown degree.

When these WMHS estimates are compared to previous global estimates, different methodological approaches and measures of these previous studies must also be taken into account—for example, whether these studies report estimates of daily tobacco smoking or daily cigarette smoking. Accordingly, this study's estimates for smoking (in general) should be higher than reported estimates of tobacco smoking or daily smoking per se. Even so, this study's range of estimates for prevalence of current smoking in several countries (25%—32%) is not appreciably different from the 1995 global estimate of 29% for daily smoking prevalence among those aged 15 and older. In addition, these WMHS estimates for a majority of the countries surveyed are not too distant from the individual country estimates reported in the most recent WHO report from 2008.

Characteristics of smokers and correlates of smoking varied somewhat across the individual countries, and in relation to the World Bank's low-medium-high gradient of economic development, but a quite prominent male excess in the odds of eversmoking was found, and one exceptional finding concerns failure to quit smoking among adult women in Spain, already mentioned in relation to possible gender-targeted outreach and intervention as part of a comprehensive tobacco prevention and control approach. The male excess odds of smoking in China is remarkable; in that country, male smoking started to increase several decades ago, while smoking among females is a more recent development and occurs substantially less frequently.<sup>31</sup> China represents a enigma for the Lopez tobacco epidemic model as it might well otherwise qualify for one of the more advanced stages. 19 Otherwise, this study's multiple country-specific smoking estimates for males and females generally are in line with the four-stage model of the tobacco epidemic proposed by Lopez and colleagues. 19

In the higher-income countries, smoking prevalence across age groups did not vary greatly or when it did (eg, in New Zealand, Belgium and Germany) the proportion of young adults (aged 18–24) smoking was found to be larger than was observed for

other age groups. Initiation still occurs frequently among individuals aged 18–24 and tobacco product marketing campaigns often target young adults. <sup>32–34</sup> On the other hand, in several of the lower-income and middle-income countries, smoking was more common among the middle adulthood age group, substantiating a need for tobacco prevention and control beyond the young adult years. Indeed, Cho and colleagues recently found that among adults in South Korea there was an age-related decline for males and an age-related increase for women. <sup>35</sup> This is a pattern not investigated in the present study, which remains on the agenda for future WMHS reports, as is an intriguing hypothesis that in some places there may be "influence of a culture which discourages married women from smoking, and "liberates" divorced women from [these] cultural sanctions" that otherwise would keep them from starting to smoke. <sup>35</sup>

Regarding employment, in all but two countries under study, more than 25% of the adult workforce were current smokers. These estimates draw attention to a potential opportunity for workplace smoke-free policies and smoking cessation programmes of the type developed elsewhere. Initiatives encouraging worksites and public places to be smoke-free may also increase the interest in creating smoke-free homes. 36

The relation between smoking and education was not monotonic and varied substantially across the WMH countries. Education is often used as an indicator of socioeconomic status (SES) as it remains fairly stable during adulthood and is correlated with occupation, income and wealth. Studies in European countries indicate that different tobacco control efforts may influence the smoking rates among various educational/SES groups differently. Perhaps country patterns of the association between education/SES and smoking reflect a stage within the tobacco epidemic, as low SES groups may follow high SES groups, after a lag, in their rejection of smoking, just as they followed high SES groups, also after a lag, in the adoption of smoking. <sup>39</sup>

In general terms, within the lower-income and middle-income countries compared to the higher-income countries, a smaller proportion of the adult population has a history of smoking. Yet in Lebanon, China and the Ukraine, the estimated proportion of adults currently smoking was higher than proportions found in most higher-income countries, and in these three countries, the estimated proportion of ever-smokers who had quit was at the lower end of the observed ranges. Increased cigarette consumption is common among countries targeted by transnational tobacco companies, 40–42 and where cigarettes have become more affordable as a result of raising household income and low excise taxes. <sup>43</sup> Countries with a high economic interest at stake (history of being a substantial producer of tobacco crops) may favour short-term financial benefits over long-term health risks.

Previous studies have found individual and environmental characteristics associated with serious quit attempts and cessation, <sup>38</sup> <sup>44–46</sup> and find that increasing age helps predict quitting. <sup>44</sup> <sup>48</sup> Among a nationally representative sample of US adults, age, race, quantity of cigarettes smoked per day, income and health were independent predictors of quitting for at least 1 year; and younger age, female gender and urban residence were predictors of relapse. <sup>48</sup> In four higher-income countries taking part in the International Tobacco Control Policy Evaluation Project, despite similarities in individual characteristics predicting making a quit attempt (intention, having tried to quit previously, duration of last quit attempt, less dependent on nicotine, more negative attitudes about smoking and younger age), differences in the success of quitting were found across countries suggesting environmental and sociocultural factors may contribute to varying trends in tobacco use. <sup>49</sup> Consistent with these WMHS

#### What this study adds

▶ This study provides new multinational findings on the basic descriptive epidemiology of smoking and cessation from adults residing in seven low-income and middle-income countries and 10 higher-income countries from all regions of the world. The estimates are based on a general field survey research protocol as applied at each country or within-country site, with self-reported current and past smoking, as implemented for the WHO World Mental Health Initiative (WMH). Observed variations in current and former smoking status across the countries underscores the reason to implement FCTC provisions while also recognising a need for locally tailored interventions.

results, Jha and colleagues<sup>8</sup> have noted a time trend that favours increased quitting among smokers in higher-income countries, but not in middle-income countries. Country differences in smoking and cessation rates may be attributable to tobacco control efforts at national levels as well as those targeting individual control. In Europe, reduced smoking rates have been found in countries with more developed and comprehensive tobacco control policies that included taxes, bans and restrictions, information campaigns, health warnings and access to treatment.<sup>38</sup>

#### CONCLUSION

It has been projected that within 25 years 10 million tobaccorelated deaths will occur annually, with 70% of these deaths in lower-income and middle-income countries. It is estimated that roughly 25 million premature deaths in the first quarter of this century and about 150 million more by mid-century might be avoided if half of the active smokers were to quit. Our findings describe the characteristics of smokers in multiple countries around the world that might be used to guide tobacco control activities and smoking cessation campaigns. The results make it clear that there is a sturdy male excess in odds of starting to smoke, but otherwise a diversity of sociodemographic associations with smoking. Moreover, the male-female variations observed for starting to smoke are not present in the patterns observed for quitting once smoking starts. As we already have mentioned in the text, the observed variations may underscore the reasons for implementation of the FCTC provisions and prompt locally tailored outreach and intervention policies and programmes for tobacco prevention and control. Spain and the situation with females who smoke but haven't quit is a case in point, as is the situation with respect to workplace smoke-free policies and programmes to encourage smoking cessation in most countries studied here. A recent Cochrane review provides guidance in this respect.  $^{50}$ 

Author affiliations: <sup>1</sup>Department of Family and Community Health, University of Maryland, Baltimore, School of Nursing, Baltimore, Maryland, USA <sup>2</sup>Department of Mental Health, Bloomberg School of Public Health, Baltimore, Maryland, USA <sup>3</sup>Department of Epidemiology, Michigan State University, College of Human Medicine, East Lansing, Michigan, USA <sup>4</sup>Health Services Research Unit, Institut Municipal d'Investigació Mèdica (IMIM-Hospital del Mar), CIBER en Epidemiología y Salud Pública (CIBERESP), Spain <sup>5</sup>Center for Public Mental Health, Gösing am Wagram, Austria <sup>6</sup>Department of Neurosciences and Psychiatry, University Hospitals Gasthuisberg, Leuven, Belgium <sup>7</sup>IRCCS Fatebenefratelli, Brescia, Italy <sup>8</sup>Netherlands Institute of Mental Health and Addiction, Utrecht, Netherlands <sup>9</sup>Department of Psychiatry, University College Hospital, Ibadan, Nigeria <sup>10</sup>Department of Psychiatry and Clinical Psychology, Balamand University School of Medicine, St George University Medical Center, Lebanon <sup>11</sup>Institute for Development, Research, Advocacy and Applied Care (IDRAAC), Beirut,

Lebanon <sup>12</sup>Ukrainian Psychiatric Association, Kiev, Ukraine <sup>13</sup>Department of Psychiatry, The Chinese University of Hong Kong, HKSAR <sup>14</sup>Hospital Ferdinand Widal, Paris, France <sup>15</sup>Department of Epidemiology, National Institute of Psychiatry, Mexico City, Mexico <sup>16</sup>School of Public Health and Family Medicine, University of Cape Town, Cape Town, South Africa <sup>17</sup>Legacy Heritage International MPH Program, Braun School of Public Health and Community Medicine, Hebrew University-Hadassah, Jerusalem, Israel <sup>18</sup>Ministry of Social Protection, Colegio Mayor de Cundinamarca University, Bogota, Colombia <sup>19</sup>Department of Preventive Cardiology, National Cardiovascular Center, Japan <sup>20</sup>Christchurch School of Medicine and Health Science, Christchurch, New Zealand <sup>21</sup>Department of Health Care Policy, Harvard Medical School, Boston, Massachusetts, USA

**Acknowledgements** The surveys discussed in this article was carried out in conjunction with the WHO World Mental Health (WMH) Survey Initiative. We thank the WMH staff for assistance with instrumentation, fieldwork and data analysis. These activities were supported by the US National Institute of Mental Health (R01MH070884), the John D and Catherine T MacArthur Foundation, the Pfizer Foundation, the US Public Health Service (R13-MH066849, R01-MH069864 and R01 DA016558), the Fogarty International Center (FIRCA R01-TW006481), the Pan American Health Organisation, Eli Lilly and Company, Ortho-McNeil Pharmaceutical, Inc. GlaxoSmithKline and Bristol-Myers Squibb. A complete list of WMH publications can be found at http://www.hcp.med.harvard.edu/wmh/. The Chinese World Mental Health Survey Initiative is supported by the Pfizer Foundation. The Colombian National Study of Mental Health (NSMH) is supported by the Ministry of Social Protection. The ESEMeD project is funded by the European Commission (contracts QLG5-1999-01042; SANCO 2004123), the Piedmont Region (Italy), Fondo de Investigación Sanitaria, Instituto de Salud Carlos III, Spain (FIS 00/0028), Ministerio de Ciencia y Tecnología, Spain (SAF 2000-158-CE), Departament de Salut, Generalitat de Catalunya, Spain, Instituto de Salud Carlos III (CIBER CB06/02/0046, RETICS RD06/ 0011 REM-TAP) and other local agencies and by an unrestricted educational grant from GlaxoSmithKline. The Israel National Health Survey is funded by the Ministry of Health with support from the Israel National Institute for Health Policy and Health Services Research and the National Insurance Institute of Israel. The World Mental Health Japan (WMHJ) Survey is supported by the Grant for Research on Psychiatric and Neurological Diseases and Mental Health (H13-SH0GAI-023, H14-T0KUBETSU-026, H16-K0K0R0-013) from the Japan Ministry of Health, Labour and Welfare. The Lebanese National Mental Health Survey (LEBANON) is supported by the Lebanese Ministry of Public Health, the WHO (Lebanon), anonymous private donations to IDRAAC, Lebanon, and unrestricted grants from Janssen Cilag, Eli Lilly, GlaxoSmithKline, Roche and Novartis. The Mexican National Comorbidity Survey (MNCS) is supported by the National Institute of Psychiatry Ramon de la Fuente (INPREMDIES 4280) and by the National Council on Science and Technology (CONACyT-G30544- H), with supplemental support from the PanAmerican Health Organisation (PAHO). Te Rau Hinengaro: The New Zealand Mental Health Survey (NZMHS) is supported by the New Zealand Ministry of Health, Alcohol Advisory Council, and the Health Research Council. The Nigerian Survey of Mental Health and Wellbeing (NSMHW) is supported by WHO (Geneva), WHO (Nigeria), and the Federal Ministry of Health, Abuja, Nigeria. The South Africa Stress and Health Study (SASH) is supported by the US National Institute of Mental Health (R01-MH059575) and National Institute of Drug Abuse with supplemental funding from the South African Department of Health and the University of Michigan. The Ukraine Comorbid Mental Disorders during Periods of Social Disruption (CMDPSD) study is funded by the US National Institute of Mental Health (RO1-MH61905. The US National Comorbidity Survey Replication (NCS-R) is supported by the National Institute of Mental Health (NIMH; U01-MH60220) with supplemental support from the National Institute of Drug Abuse (NIDA), the Substance Abuse and Mental Health Services Administration (SAMHSA), the Robert Wood Johnson Foundation (RWJF; Grant 044708) and the John W Alden Trust.

**Contributors** CLS conceptualised the idea and wrote the manuscript. HC performed the data analysis. JCA provided oversight, supervised the analyses, and was involved in revisions of the original manuscript. All other authors were involved in the WMH collaborative effort that provided the data and provided comments on the manuscript.

Funding The preparation of this manuscript was funded by the US National Institute on Drug Abuse (R01DA016558 and K05DA015799). The surveys discussed in this article was carried out in conjunction with the World Health Organization World Mental Health (WMH) Survey Initiative. We thank the WMH staff for assistance with instrumentation, fieldwork, and data analysis. These activities were supported by the United States National Institute of Mental Health (R01MH070884), the John D and Catherine T MacArthur Foundation, the Pfizer Foundation, the US Public Health Service (R13-MH066849, R01-MH069864, and R01 DA016558), the Fogarty International Center (FIRCA R01-TW006481), the Pan American Health Organization, Eli Lilly and Company, Ortho-McNeil Pharmaceutical, Inc, GlaxoSmithKline, and Bristol-Myers Squibb. A complete list of WMH publications can be found at http://www.hcp.med.harvard.edu/wmh/. The Chinese World Mental Health Survey Initiative is supported by the Pfizer Foundation. The Colombian National Study of Mental Health (NSMH) is supported by the Ministry of Social Protection. The ESEMeD project is funded by the European Commission (Contracts QLG5-1999-01042; SANCO 2004123), the Piedmont

Region (Italy), Fondo de Investigación Sanitaria, Instituto de Salud Carlos III, Spain (FIS 00/0028), Ministerio de Ciencia y Tecnología, Spain (SAF 2000-158-CE), Departament de Salut, Generalitat de Catalunya, Spain, Instituto de Salud Carlos III (CIBER CB06/02/0046, RETICS RD06/0011 REM-TAP), and other local agencies and by an unrestricted educational grant from GlaxoSmithKline. The Israel National Health Survey is funded by the Ministry of Health with support from the Israel National Institute for Health Policy and Health Services Research and the National Insurance Institute of Israel. The World Mental Health Japan (WMHJ) Survey is supported by the Grant for Research on Psychiatric and Neurological Diseases and Mental Health (H13-SHOGAI-023, H14-TOKUBETSU-026, H16-KOKORO-013) from the Japan Ministry of Health, Labour and Welfare, The Lebanese National Mental Health Survey (LEBANON) is supported by the Lebanese Ministry of Public Health, the WHO (Lebanon), anonymous private donations to IDRAAC, Lebanon, and unrestricted grants from Janssen Cilag, Eli Lilly, GlaxoSmithKline, Roche, and Novartis. The Mexican National Comorbidity Survey (MNCS) is supported by The National Institute of Psychiatry Ramon de la Euente (INPREMDIES 4280) and by the National Council on Science and Technology (CONACyT-G30544- H), with supplemental support from the PanAmerican Health Organization (PAHO). Te Rau Hinengaro: The New Zealand Mental Health Survey (NZMHS) is supported by the New Zealand Ministry of Health, Alcohol Advisory Council, and the Health Research Council. The Nigerian Survey of Mental Health and Wellbeing (NSMHW) is supported by the WHO (Geneva), the WHO (Nigeria), and the Federal Ministry of Health, Abuja, Nigeria. The South Africa Stress and Health Study (SASH) is supported by the US National Institute of Mental Health (R01-MH059575) and National Institute of Drug Abuse with supplemental funding from the South African Department of Health and the University of Michigan. The Ukraine Comorbid Mental Disorders during Periods of Social Disruption (CMDPSD) study is funded by the US National Institute of Mental Health (RO1-MH61905). The US National Comorbidity Survey Replication (NCS-R) is supported by the National Institute of Mental Health (NIMH; U01-MH60220) with supplemental support from the National Institute of Drug Abuse (NIDA), the Substance Abuse and Mental Health Services Administration (SAMHSA), the Robert Wood Johnson Foundation (RWJF; Grant 044708), and the John W Alden Trust. Other funders:NIH.

#### Conflict of interests None.

**Ethics approval** The institutional review board of the organisation coordinating the survey in each country approved and monitored the compliance of human subject protection and obtaining informed consent. Data analysis activities were approved by the Johns Hopkins Bloomberg School of Public Health and Michigan State University institutional review boards.

Provenance and peer review Not commissioned; externally peer reviewed.

#### **REFERENCES**

- Corraro MA, Guindon GE, Sharma N, et al, eds. Tobacco control country profiles. Atlanta, GA: American Cancer Society, 2000.
- Global Youth Tobacco Survey Collaborative Group. Tobacco use among youth: a cross country comparison. Tob Control 2002;11:252—70.
- Shafey 0, Dolwixk S, Guindon GE, eds. Tobacco control country profiles. 2nd ed. Atlanta. GA: American Cancer Society, 2003.
- Warren CW, Jones NR, Eriksen MP, et al. Patterns of global tobacco use in young people and implications for future chronic disease burden in adults. Lancet 2006;367:749—53.
- Ezzati M, Henley SJ, Lopez AD, et al. Role of smoking in global and regional cancer epidemiology:current patterns and data needs. Int J Cancer 2005;116:963—71.
- World Health Organization. Tobacco or health:a global status report. Geneva: World Health Organization, 1997.
- Peto RL, Lopez AD. The future worldwide health effects of current smoking patterns. In: Koop CE, Pearson CE, Schwarz MR, eds. *Critical issues in global health*. San Francisco, CA: Jossey-Bass, 2000.
- Jha P, Ranson MK, Nguyen SN, et al. Estimates of global and regional smoking prevalence in 1995, by age and sex. Am J Public Health 2002;92:1002—6.
- WHO report on the global tobacco epidemic, 2008:the MPOWER package. Geneva: World Health Organization 2008.
- Fagerström KO, Kunze M, Schoberberger R, et al. Nicotine dependence versus smoking prevalence:comparisons among countries and categories of smokers. Tob Control 1996;5:52—6.
- Fagerström K, Furberg H. A comparison of the Fagerström test for nicotine dependence and smoking prevalence across countries. *Addiction* 2008;103:841-5.
- Dormitzer CM, Gonzalez GB, Penna M, et al. The PACARDO research project: youthful drug involvement in Central America and the Dominican Republic. Rev Panam Salud Publica 2004;15:400—16.
- ESPAD. European school survey project on alcohol and other drugs. http://www. espad.org/espad-reports (accessed 26 May 2009).
- GATS. Global adult tobacco survey. http://www.searo.who.int/EN/Section1174/ Section2469/Section2482 14011.htm (accessed 26 May 2009).
- World Health Survey. http://www.who.int/healthinfo/survey/en/index.html (accessed 26 May 2009).
- STEPwise approach to chronic disease risk factor surveillance (STEPS). http://www. who.int/chp/steps/riskfactor/en/index.html (accessed 26 May 2009).

- ITC Project. The International Tobacco Control Policy Evaluation Project. http://www.itcproject.org/ (accessed 18 Sep 2009).
- Fong GT, Cummings KM, Shopland DR. Building the evidence base for effective tobacco control policies: the International Tobacco Control Policy Evaluation Project (the ITC Project). Tob Control 2006;15(suppl 3):iii1—iii2.
- Lopez AD, Collishaw NE, Piha T. A descriptive model of the cigarette epidemic in developed countries. *Tob Control* 1994;3:242—7.
- MacKay J, Ericksen M. The tobacco atlas. Geneva: World Health Organization, 2002
- United States Department of Health and Human Services (US DHHS), Public Health Service. The health benefits of smoking cessation: a report of the Surgeon General. Washington, DC: US Government Printing Office, DHHS, 1990 publication no. (CDC) 90-8416.
- Dresler CM, Leon ME, Straif K, et al. Reversal of risk upon quitting smoking. Lancet 2006:368:348—9.
- Peto R, Darby S, Deo S, et al. Smoking, smoking cessation, and lung cancer in the UK since 1950: combination of national statistics with two case-control studies. BMJ 2000;321:323—9.
- Demyttenaere K, Bruffaerts R, Posada-Villa J, et al. Prevalence, severity, and unmet need for treatment of mental disorders in the World Health Organization World Mental Health Surveys. JAMA 2004;291:2581—90.
- Kessler RC, Ustun TB. The World Mental Health (WMH) Survey Initiative Version of the World Health Organization (WHO) Composite International Diagnostic Interview (CIDI). Int J Methods Psychiatr Res 2004;13:93—121.
- Kessler RC, Üstün TB, eds. The WHO World Mental Health Survey:global perspectives on the epidemiology of mental disorders. Cambridge: Cambridge University Press, 2008.
- World Bank. World trade indicators. 2008. http://info.worldbank.org/etools/wti2008/ docs/Countrylist.htm (accessed 30 Jun 2008).
- Storr CL, Chen H, Posada-Villa J, et al. Adult smokers in Columbia: who isn't giving it up? Addict Behav 2008:33:412—21.
- Stata Corporation. Stata statistical software:release 9. College Station, TX: Stata Corporation, 2005.
- Borland R, Cummings KM, Leon M, et al. Methods for evaluating tobacco control policies. International Agency for Research on Cancer (IARC), handbook 12 on cancer prevention, tobacco control. Geneva: World Health Organization, 2009
- Gong YL, Koplan JP, Feng W, et al. Cigarette smoking in China. Prevalence, characteristics, and attitudes in Minhang District. JAMA 1995;274:1232–4.
- Sargent JD, Hanewinkel R. Comparing the effects of entertainment media and tobacco marketing on youth smoking in Germany. Addiction 2009;104:815—23.
- Cortese DK, Lewis MJ, Ling PM. Tobacco industry lifestyle magazines targeted to young adults. J Adolesc Health 2009;45:268—80.
- Sepe E, Ling PM, Glantz SA. Smooth moves:bar and nightclub tobacco promotions that target young adults. Am J Public Health 2002;92:414—19.
- Cho HJ, Khang YH, Jun HJ, et al. Marital status and smoking in Korea: the influence of gender and age. Soc Sci Med 2008;66:609—19.
- Borland R, Yong HH, Cummings KM, et al. Determinants and consequences of smoke-free homes: findings from the International Tobacco Control (ITC) Four Country Survey. Tob Control 2006;15(Suppl 3):iii42—50.
- Giskes K, Kunst AE, Benach J, et al. Trends in smoking behaviour between 1985 and 2000 in nine European countries by education. J Epidemiol Community Health 2005;59:395—401.
- Schaap MM, Kunst AE, Leinsalu M, et al. Effect of nationwide tobacco control policies on smoking cessation in high and low educated groups in 18 European countries. Tob Control 2008;17:248—55.
- Cavelaars A, Kunst A, Geurtts J, et al. Educational differences in smoking: international comparison. BMJ 2000;320:1102—7.
- Nakkash R, Lee K. Smuggling as the "key to a combined market": British American Tobacco in Lebanon. *Tob Control* 2008;17:324—31.
- Nakkash R, Lee K. The tobacco industry's thwarting of marketing restrictions and health warnings in Lebanon. *Tob Control* 2009;18:310—16.
- Zhong F, Yano E. British American tobacco's tactics during China's accession to the World Trade Organization. *Tob Control* 2007;16:133—7.
- Peng L, Ross H. The impact of cigarette taxes and advertising on the demand for cigarettes in Ukraine. Cent Eur J Public Health 2009;17:93—8.
- 44. **Hatziandreu EJ**, Pierce JP, Lefkopoulou M, *et al.* Quitting smoking in the United States in 1986. *J Natl Cancer Inst* 1990;**82**:1402—6.
- Tucker JS, Ellickson PL, Orlando M, et al. Predictors of attempted quitting and cessation among young adult smokers. Prev Med 2005;41:554—61.
- Giskes K, van Lenthe FJ, Turrell G, et al. Smokers living in deprived areas are less likely to quit: a longitudinal follow-up. Tob Control 2006;15:485—8.
- Hyland A, Li Q, Bauer JE, et al. Predictors of cessation in a cohort of current and former smokers followed over 13 years. Nicotine Tob Res 2004;6(suppl 3): \$363\_0
- 48. **McWhorter WP**, Boyd GM, Mattson ME. Predictors of quitting smoking: the NHANES I followup experience. *J Clin Epidemiol* 1990; **43**:1399—405.
- Hyland A, Borland R, Li Q, et al. Individual-level predictors of cessation behaviours among participants in the International Tobacco Control (ITC) Four Country Survey. Tob Control 2006;15(suppl 3):iii83—iii94.
- Cahill K, Moher M, Lancaster T. Workplace interventions for smoking cessation. Cochrane Database Syst Rev 2008;4:CD003440.