



Tables from the Dartmouth Atlas Ambulatory Care Quality Report: State Level

State name	Major lower extremity amputations per 1,000 Medicare enrollees (2003-05)			Percent of female Medicare enrollees age 65-69 having at least one mammogram over the two-year period (2004-2005)			Average annual percent of Medicare enrollees with diabetes age 65-74 having hemoglobin A1c test (2003-05)			Percent of Medicare beneficiaries whose predominant ambulatory provider was a primary care physician (2004)			Discharges for ambulatory care sensitive conditions per 1,000 Medicare enrollees (2003-05)		
	<i>Whether patients face amputation, a devastating complication of peripheral vascular disease and diabetes</i>			<i>Whether women get basic recommended care, such as mammography tests</i>			<i>Whether patients with diabetes are getting recommended blood tests</i>			<i>The primary care orientation of their local care system</i>			<i>How often patients are admitted to hospitals because their chronic conditions like diabetes or heart failure are poorly managed</i>		
	Overall	Black	White	Overall	Black	White	Overall	Black	White	Overall	Black	White	Overall	Black	White
Alabama	1.49	5.37	1.16	63.5	58.8	64.4	82.1	79.3	83.1	81.5	85.3	81.0	96.5	106.6	98.3
Alaska	0.90			58.3			70.9			76.7			63.4	48.6	62.4
Arizona	0.85	2.63	0.67	64.6			77.4	80.9	77.3	75.7	78.7	75.5	59.3	76.0	57.6
Arkansas	1.35	5.38	1.00	60.0	52.1	60.8	81.1	76.9	81.8	83.2	85.0	83.0	94.7	108.0	94.0
California	0.99	3.30	0.78	59.3	47.8	60.0	80.0	74.3	80.5	72.4	74.9	72.5	64.3	100.1	61.7
Colorado	0.67	1.76	0.53	62.7			83.0	66.1	84.0	80.2	79.2	80.0	56.1	65.8	54.7
Connecticut	0.96	2.88	0.78	68.8	63.8	69.3	86.5	83.0	87.0	69.8	72.4	69.6	65.3	80.6	63.7
Delaware	1.05	3.80	0.81	69.4	66.5	69.8	84.1	79.8	85.2	75.0	80.6	74.4	73.3	99.8	70.9
District of Columbia	0.90	3.39	0.47	58.9	55.6	65.5	81.3	80.8	84.6	60.2	63.2	58.0	64.1	93.4	46.6
Florida	0.98	4.10	0.73	67.6	58.6	68.3	84.4	81.5	84.9	71.3	79.2	70.8	69.8	96.0	67.6
Georgia	1.37	5.21	1.00	63.3	57.3	64.9	83.5	81.4	84.3	74.8	79.0	74.2	83.1	94.2	84.8
Hawaii	0.92			57.3			87.3			79.5			31.8	40.2	30.9
Idaho	0.71			61.1			85.6			75.2			55.5	75.1	53.9
Illinois	1.05	3.32	0.86	61.6	51.0	63.0	81.7	70.1	84.0	77.1	78.7	77.0	87.7	129.9	83.9
Indiana	1.13	3.49	0.91	61.9	56.3	62.3	83.1	74.3	84.0	81.0	81.2	80.8	81.8	101.7	79.9
Iowa	0.86	3.03	0.67	67.3			89.4	87.1	89.5	83.3	85.0	83.0	67.8	93.0	65.8
Kansas	0.89	2.42	0.72	66.0			86.5	76.4	87.2	84.6	85.5	84.4	79.6	89.2	77.8
Kentucky	1.27	3.75	1.03	61.9	65.2	61.7	83.4	84.8	83.3	82.0	85.1	81.7	112.9	104.5	112.1
Louisiana	1.66	6.14	1.26	60.9	56.7	62.1	79.3	77.1	80.3	72.0	78.3	70.7	109.9	130.4	111.4
Maine	0.98			74.0			90.8			85.3			67.3	67.6	65.4
Maryland	1.00	3.55	0.79	64.1	60.4	65.1	84.8	80.7	86.6	74.5	76.6	74.3	75.2	99.6	73.2
Massachusetts	0.93	2.30	0.76	71.9	71.5	71.9	89.0	87.8	89.1	74.3	82.2	74.0	80.1	96.0	78.2
Michigan	0.91	2.99	0.74	69.8	62.6	70.7	86.1	80.6	87.1	81.8	82.8	81.7	76.9	111.7	73.8
Minnesota	0.88	2.19	0.69	70.7			90.2	83.2	90.3	83.9	82.0	83.6	64.5	81.2	62.7
Mississippi	1.60	5.61	1.31	56.9	49.3	59.5	77.8	76.4	78.6	76.6	83.0	75.1	109.1	127.9	112.1

Blank cells indicate that there was an insufficient number of black patients in the category to report race-specific rates.



Tables from the Dartmouth Atlas Ambulatory Care Quality Report: State Level

State name	Major lower extremity amputations per 1,000 Medicare enrollees (2003-05)			Percent of female Medicare enrollees age 65-69 having at least one mammogram over the two-year period (2004-2005)			Average annual percent of Medicare enrollees with diabetes age 65-74 having hemoglobin A1c test (2003-05)			Percent of Medicare beneficiaries whose predominant ambulatory provider was a primary care physician (2004)			Discharges for ambulatory care sensitive conditions per 1,000 Medicare enrollees (2003-05)		
	<i>Whether patients face amputation, a devastating complication of peripheral vascular disease and diabetes</i>			<i>Whether women get basic recommended care, such as mammography tests</i>			<i>Whether patients with diabetes are getting recommended blood tests</i>			<i>The primary care orientation of their local care system</i>			<i>How often patients are admitted to hospitals because their chronic conditions like diabetes or heart failure are poorly managed</i>		
	Overall	Black	White	Overall	Black	White	Overall	Black	White	Overall	Black	White	Overall	Black	White
Missouri	1.14	4.19	0.88	62.0	59.0	62.3	84.3	78.1	85.1	82.3	81.1	82.3	87.1	112.8	84.8
Montana	0.82			70.0			82.7			80.5			68.9	63.5	67.0
Nebraska	0.85	2.24	0.67	63.3			87.2	82.7	87.3	86.4	81.8	86.3	70.3	89.6	68.3
Nevada	0.71	1.50	0.60	57.9			78.3	77.1	78.4	72.6	73.2	72.5	63.5	76.3	62.1
New Hampshire	0.96			70.6			89.3			81.0			62.7	58.8	61.0
New Jersey	1.06	3.73	0.83	59.1	54.2	59.7	80.9	74.6	82.1	65.3	71.8	64.8	83.9	125.2	80.3
New Mexico	0.96			58.8			73.6			80.0	82.6	80.0	64.9	66.5	63.3
New York	1.08	3.47	0.88	60.7	51.6	61.8	83.6	76.8	84.8	70.5	71.6	70.4	78.7	105.6	76.3
North Carolina	1.40	5.31	1.04	67.1	60.1	68.5	87.4	86.1	87.9	81.5	84.8	81.1	76.0	94.0	75.3
North Dakota	1.03			73.5			88.7			82.6			71.5		
Ohio	1.08	3.31	0.89	65.4	62.0	65.7	84.7	80.1	85.3	82.1	83.2	82.0	87.3	109.0	85.4
Oklahoma	1.20	3.64	0.96	57.1	53.9	57.2	77.6	77.2	77.6	81.3	80.2	81.3	95.4	100.3	93.8
Oregon	0.85			66.1			87.7			83.3	85.1	83.0	52.3	62.0	50.8
Pennsylvania	1.29	3.35	1.07	64.0	53.7	64.6	85.5	79.1	86.0	82.1	81.3	82.0	84.6	121.7	81.7
Rhode Island	0.94			63.7			84.4			78.7	83.7	78.4	77.8	101.9	75.6
South Carolina	1.56	6.04	1.12	65.3	60.3	66.6	82.0	79.9	83.1	80.0	83.9	79.4	77.3	94.3	77.5
South Dakota	1.06			65.3			85.4			86.4			76.0	53.0	73.8
Tennessee	1.36	4.57	1.09	62.7	57.8	63.2	86.8	81.4	87.8	81.0	82.5	80.9	98.6	107.6	98.6
Texas	1.50	4.20	1.27	58.7	54.0	59.1	82.9	80.7	83.2	78.3	82.5	78.1	88.6	105.8	87.2
Utah	0.50			61.1			86.5			81.1			47.6	59.4	46.3
Vermont	1.02			69.6			91.5			82.5			58.7	65.7	57.0
Virginia	1.12	4.11	0.86	63.6	59.2	64.5	85.8	81.9	87.3	78.9	81.9	78.5	71.2	84.2	71.2
Washington	0.92	2.54	0.72	65.5			89.1	81.6	89.3	79.6	77.6	79.5	50.0	64.5	48.6
West Virginia	1.36	3.88	1.08	64.4			83.8	78.3	84.0	80.9	79.9	80.7	116.4	124.0	113.8
Wisconsin	1.20	4.09	0.93	70.2	61.3	70.5	90.9	85.4	91.2	81.6	80.0	81.4	63.2	98.7	61.1
Wyoming	0.91			60.9			78.1			79.7			68.1	54.1	66.3
United States	1.14	4.17	0.88	63.6	56.8	64.3	84.0	79.4	84.7	77.6	80.0	77.4	78.3	105.0	76.0

Blank cells indicate that there was an insufficient number of black patients in the category to report race-specific rates.